

Flavor nonconservation and CP violation from quark mixings with singlet quarks

Isao Kakebe ¹, Katsuji Yamamoto ²

Department of Nuclear Engineering, Kyoto University, Kyoto 606-01, Japan

abstract

Flavor nonconserving and CP violating effects of the quark mixings are investigated in electroweak models incorporating singlet quarks. Especially, the D^0 - $\bar{D}^0$ mixing and the neutron electric dipole moment, which are mediated by the up-type quark couplings to the neutral Higgs fields, are examined in detail for reasonable ranges of the quark mixings and the singlet Higgs mass scale. These neutral Higgs contributions are found to be comparable to or smaller than the experimental bounds even for the case where the singlet Higgs mass scale is of the order of the electroweak scale and a significant mixing is present between the top quark and the singlet quarks.

¹ e-mail address: kakebe@b1.nucleng.kyoto-u.ac.jp

² e-mail address: yamamoto@b1.nucleng.kyoto-u.ac.jp

Some extensions of the minimal standard model may be motivated in various points of view. For instance, in the electroweak baryogenesis, the CP asymmetry induced by the conventional phase in the Cabibbo-Kobayashi-Maskawa (CKM) matrix is far too small to account for the observed baryon to entropy ratio, and the electroweak phase transition should be at most of weakly first order consistently with the experimental bound on the Higgs particle mass [1]. In this article, among various possible extensions, we investigate electroweak models incorporating $SU(2)_W \times U(1)_Y$ singlet quarks with electric charges $2/3$ and/or $-1/3$ [2, 3, 4, 5, 6, 7, 8, 9, 10, 11]. These sorts of models have some novel features arising from the mixings between the ordinary quarks (q) and the singlet quarks (Q): The CKM unitarity in the ordinary quark sector is violated, and the flavor changing neutral currents (FCNC's) are present at the tree-level. Furthermore, the q - Q mixings involving new CP violating sources are expected to make important contributions to the electroweak baryogenesis [12]. A singlet Higgs field S introduced to provide the singlet quark mass terms and q - Q mixing terms is even preferable for realizing a strong enough first order electroweak phase transition [1]. It should also be mentioned, as investigated in the earlier literature [6], that the complex vacuum expectation value (vev) $\langle S \rangle$ of the singlet Higgs field induced by the spontaneous CP violation leads to a non-vanishing CKM phase.

The quark mixings with singlet quarks are, on the other hand, subject to the constraints coming from various flavor nonconserving and CP violating processes [2, 3, 4, 5, 6, 8, 10, 11]. In particular, it is claimed in [6] with a simple calculation for the case of d - D mixings that a rather stringent bound, $v_S \equiv |\langle S \rangle| \gtrsim$ several TeV, on the singlet Higgs mass scale is imposed from the one-loop neutral Higgs contributions to the neutron electric dipole moment (NEDM). (The NEDM in electroweak models with singlet quarks is also considered in [3, 4].) The singlet Higgs field with mass scale in TeV region, however, seems to be unfavorable for the electroweak baryogenesis. Considering this apparently controversial situation, we examine in detail such phenomenological implications of the electroweak models with singlet quarks. We mainly describe the case of singlet U quarks with electric charge $2/3$ in the following, keeping in mind the possibility of significant t - U mixing for the electroweak baryogenesis. The analyses are extended readily to the case of singlet D quarks with electric charge $-1/3$, and similar results are obtained for both cases. In practice, diagonalization of the quark mass matrix is made numerically, in order to calculate precisely

the relevant quark couplings to the neutral Higgs fields involving flavor nonconservation and CP violation. Then, for reasonable ranges of the model parameters, systematic analyses are performed on the neutral Higgs contributions to the D^0 - $\bar{D}^0$ mixing and the NEDM. In contrast to the earlier expectation [6], they will turn out to be comparable to or smaller than the experimental bounds even for the case where the singlet Higgs mass scale is of the order of the electroweak scale and a significant t - U mixing is present.

The Yukawa couplings relevant for the up-type quark sector are given by

$$\mathcal{L}_{\text{Yukawa}} = -u_0^c \lambda_u q_0 H - u_0^c (f_U S + f'_U S^\dagger) U_0 - U_0^c (\lambda_U S + \lambda'_U S^\dagger) U_0 + \text{h.c.} \quad (1)$$

with the two-component Weyl fields (the generation indices and the factors representing the Lorentz covariance are omitted for simplicity). Here $q_0 = (u_0, V_0 d_0)$ represents the quark doublets with a unitary matrix V_0 , and $H = (H^0, H^+)$ is the electroweak Higgs doublet. A suitable redefinition among the u_0^c and U_0^c fields with the same quantum numbers has been made to eliminate the $U_0^c q_0 H$ couplings without loss of generality. Then, the Yukawa coupling matrix λ_u has been made diagonal by using unitary transformations among the ordinary quark fields. In this basis, by turning off the u - U mixings with $f_U, f'_U \rightarrow 0$, u_0 and d_0 are reduced to the mass eigenstates, and V_0 is identified with the CKM matrix. The actual CKM matrix is slightly modified due to the u - U mixings (and possibly the d - D mixings), as shown explicitly later. The Higgs fields develop vev's,

$$\langle H^0 \rangle = \frac{v}{\sqrt{2}}, \quad \langle S \rangle = e^{i\phi_S} \frac{v_S}{\sqrt{2}}, \quad (2)$$

where $\langle S \rangle$ may acquire a nonvanishing phase ϕ_S due to either spontaneous or explicit CP violation originating in the Higgs sector. The quark mass matrix is produced with these vev's as

$$\mathcal{M}_U = \begin{pmatrix} M_u & \Delta_{u-U} \\ 0 & M_U \end{pmatrix}, \quad (3)$$

where

$$M_u = \frac{\lambda_u v}{\sqrt{2}}, \quad \Delta_{u-U} = (f_U e^{i\phi_S} + f'_U e^{-i\phi_S}) \frac{v_S}{\sqrt{2}}, \quad M_U = (\lambda_U e^{i\phi_S} + \lambda'_U e^{-i\phi_S}) \frac{v_S}{\sqrt{2}}. \quad (4)$$

The quark mass matrix is diagonalized by unitary transformations $\mathcal{V}_{U_L}$ and $\mathcal{V}_{U_R}$ as

$$\mathcal{V}_{U_R}^\dagger \mathcal{M}_U \mathcal{V}_{U_L} = \text{diag.}(m_u, m_c, m_t, m_U). \quad (5)$$

(While the case with one singlet U quark is described hereafter for simplicity of notation, the analyses are readily extended to the case with more than one singlet U quarks, resulting analogous conclusions.) The quark mass eigenstates are determined in terms of the original states by

$$\begin{pmatrix} u \\ U \end{pmatrix} = \mathcal{V}_{\mathcal{U}_L}^\dagger \begin{pmatrix} u_0 \\ U_0 \end{pmatrix}, \quad (u^c, U^c) = (u_0^c, U_0^c) \mathcal{V}_{\mathcal{U}_R}. \quad (6)$$

The unitary transformations for the quark mixings may be represented as

$$\mathcal{V}_{\mathcal{U}_\chi} = \begin{pmatrix} V_{u_\chi} & \epsilon_{u_\chi} \\ -\epsilon_{u_\chi}^\dagger & V_{U_\chi} \end{pmatrix} \quad [\chi = \text{L, R}]. \quad (7)$$

(Similar unitary transformations $\mathcal{V}_{\mathcal{D}_\chi}$ are introduced if the d - D mixings are also present.)

Then, the u - U mixing submatrices are found in the leading orders to be

$$(\epsilon_{u_L})_i \simeq (\epsilon'_{u_L})_i \simeq (M_u \Delta_{u-U} M_U^{-2})_i \sim (m_{u_i}/m_U) \epsilon_{u-U}, \quad (8)$$

$$(\epsilon_{u_R})_i \simeq (\epsilon'_{u_R})_i \simeq (\Delta_{u-U} M_U^{-1})_i \sim \epsilon_{u-U}. \quad (9)$$

Here the parameter $\epsilon_{u-U} \sim (|f_U| + |f'_U|)/(|\lambda_U| + |\lambda'_U|)$ represents the mean magnitude of the u - U mixings, though there may be some generation dependence more precisely, as seen later. The relation between the masses and Yukawa couplings of the ordinary up-type quarks is slightly modified by the u - U mixings as

$$m_{u_i} \simeq \frac{\lambda_{u_i} v}{\sqrt{2}} \left[1 - \frac{1}{2} (\epsilon_{u_R} \epsilon_{u_R}^\dagger)_{ii} \right]. \quad (10)$$

The generalized CKM matrix for the up-type quarks including the U quark is given by

$$\begin{pmatrix} V & 0 \\ 0 & 0 \end{pmatrix} = \mathcal{V}_{\mathcal{U}_L}^\dagger \begin{pmatrix} V_0 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} V_{u_L}^\dagger V_0 & 0 \\ \epsilon_{u_L}^\dagger V_0 & 0 \end{pmatrix}. \quad (11)$$

(The effect of possible d - D mixings may be included by multiplying $\mathcal{V}_{\mathcal{D}_L}$ from the right.)

Here the CKM unitarity within the ordinary quark sector is violated slightly due to the u - U mixings.

It should be noticed in eq.(8) that the left-handed u - U mixings are suppressed further by the relevant u - U mass ratios m_{u_i}/m_U . On the other hand, the right-handed u - U mixings given in eq.(9) are actually ineffective by themselves, since u^c and U^c fields have the same quantum numbers. Hence, in models of this sort with the quark mass matrix of the form

given in eqs.(3) and (4), the effects of the light ordinary quark mixings with the singlet quarks appear to be rather small with suppression factors of ϵ_{u-U} and m_{u_i}/m_U under the natural relation $\lambda_{u_i} \sim m_{u_i}/v$. In fact, the CKM unitarity violation within the ordinary quark sector is found to arise at the order of $(m_{u_i}m_{u_j}/m_U^2)\epsilon_{u-U}^2$ [2, 3, 5, 6, 10, 11], which is sufficiently below the experimental bounds [13]. Contrary to this situation, it is in principle possible, for instance, to take $\lambda_{u_1} \gg m_u/v$ by making a fine-tuning in eq.(10) with a significant mixing between the $u(=u_1)$ quark and the singlet U quark. However, such a choice will not respect the quark mass hierarchy in a natural sense; the smallness of m_u is no longer guaranteed by a chiral symmetry appearing for $\lambda_{u_1} \rightarrow 0$.

The quark couplings to the neutral Higgs fields are extracted from the Yukawa couplings (1), which involve flavor nonconservation and CP violation induced by the $u-U$ mixings:

$$\mathcal{L}_{\text{Yukawa}}^{\text{neutral}} = - \sum_{a=0,1,2} \mathcal{U}^c \Lambda_a \mathcal{U} \phi_a + \text{h.c.} , \quad (12)$$

where $\mathcal{U} = (u, c, t, U)$ represents the quark mass eigenstates, and ϕ_0, ϕ_1, ϕ_2 are the mass eigenstates of the neutral Higgs fields. The original complex Higgs fields are decomposed as $H^0 = \langle H^0 \rangle + (h_1 + ih_2)/\sqrt{2}$ and $S = \langle S \rangle + (s_1 + is_2)/\sqrt{2}$ with real fields. While the Nambu-Goldstone mode h_2 is absorbed by the Z gauge boson, the remaining h_1, s_1, s_2 are combined to form the mass eigenstates ϕ_a . Then, the coupling matrices in eq.(12) are given by

$$\Lambda_a = \frac{1}{\sqrt{2}} \mathcal{V}_{\mathcal{U}_R}^\dagger (\mathcal{O}_{a0} \Lambda_u + \mathcal{O}_{a1} \Lambda_U^+ + i \mathcal{O}_{a2} \Lambda_U^-) \mathcal{V}_{\mathcal{U}_L} \quad (13)$$

with

$$\Lambda_u = \begin{pmatrix} \lambda_u & 0 \\ 0 & 0 \end{pmatrix} , \quad \Lambda_U^\pm = \begin{pmatrix} 0 & f_U \pm f'_U \\ 0 & \lambda_U \pm \lambda'_U \end{pmatrix} . \quad (14)$$

Here the Higgs mass eigenstates are expressed with a suitable orthogonal matrix $\mathcal{O}$ as

$$\begin{pmatrix} \phi_0 \\ \phi_1 \\ \phi_2 \end{pmatrix} = \mathcal{O} \begin{pmatrix} h_1 \\ s_1 \\ s_2 \end{pmatrix} . \quad (15)$$

At present the Higgs masses m_{ϕ_a} and the mixing matrix $\mathcal{O}$ should be regarded as free parameters varying in certain reasonable ranges.

The quark mass matrix may be diagonalized perturbatively with respect to the relevant couplings to determine the quark mixing matrices $\mathcal{V}_{\mathcal{U}_L}$ and $\mathcal{V}_{\mathcal{U}_R}$. Then, it is seen from eqs.(13)

and (14) that the FCNC's of the ordinary up-type quarks coupled to the neutral Higgs fields, in particular, have the following specific generation dependence:

$$(\Lambda_a)_{ij} \sim (m_{u_j}/m_U)\epsilon_{u-U}^2 \quad (i \neq j) . \quad (16)$$

This feature is actually confirmed by numerical calculations, and is also valid for the case of d - D mixings. Since these FCNC's coupled to the neutral Higgs fields are of the first order of the ordinary quark Yukawa couplings, their contributions are expected to be significant in certain flavor nonconserving and CP violating processes such as the NEDM, D^0 - $\bar{D}^0$, K^0 - $\bar{K}^0$, B^0 - $\bar{B}^0$, $b \rightarrow s\gamma$, and so on [2, 3, 4, 5, 6, 8, 10, 11]. In fact it will be seen below that the neutral Higgs contributions to the D^0 - $\bar{D}^0$ mixing and the NEDM become important for reasonable ranges of the model parameters. In contrast to the FCNC's coupled to the neutral Higgs fields, the q - Q mixing effects on the Z gauge boson couplings appear at the order of $(m_{u_i}m_{u_j}/m_U^2)\epsilon_{u-U}^2$ with the relation $m_{u_i} \sim \lambda_{u_i}/v$, which are related to the CKM unitarity violation [2, 3, 5, 10, 11]. Since they are of the second order of the u - U mass ratios, the contributions of the FCNC's in gauge couplings are suppressed much more, being sufficiently below the experimental bounds [13] for the natural choices of the model parameters. Detailed analyses on various flavor nonconserving and CP violating effects coming from the quark mixings with singlet quarks will be presented elsewhere, which are, in particular, mediated by the quark couplings to the neutral Higgs fields. They would serve as signals for the new physics beyond the minimal standard model.

The effective Hamiltonian relevant for the D^0 - $\bar{D}^0$ mixing is obtained with the quark couplings in eq. (13) mediated by the neutral Higgs fields ϕ_a . They are written with the four-component Dirac fields as

$$\mathcal{H}_\phi^{\Delta c=2} = \sum_a \frac{1}{m_{\phi_a}^2} \left[\bar{c} \left\{ (\Gamma_a^S)_{21} + (\Gamma_a^P)_{21} \gamma_5 \right\} u \right]^2 , \quad (17)$$

where

$$(\Gamma_a^S)_{21} = \frac{1}{2} [(\Lambda_a)_{12}^* + (\Lambda_a)_{21}] , \quad (\Gamma_a^P)_{21} = \frac{1}{2} [(\Lambda_a)_{12}^* - (\Lambda_a)_{21}] . \quad (18)$$

The D^0 - $\bar{D}^0$ transition matrix element is calculated with this effective Hamiltonian as follows [14]:

$$\begin{aligned} \langle \bar{D}^0 | \mathcal{H}_\phi^{\Delta c=2} | D^0 \rangle &= \sum_a \frac{1}{6m_{\phi_a}^2} \left[\left\{ (\Gamma_a^S)_{21}^2 - (\Gamma_a^P)_{21}^2 \right\} \mathcal{M}_A^0 \right. \\ &\quad \left. + \left\{ -(\Gamma_a^S)_{21}^2 + 11(\Gamma_a^P)_{21}^2 \right\} \mathcal{M}_P^0 \right] , \end{aligned} \quad (19)$$

$$\mathcal{M}_A^0 = f_D^2 m_D^2, \quad \mathcal{M}_P^0 = \frac{f_D^2 m_D^4}{(m_u + m_c)^2}. \quad (20)$$

(In the present analysis, it is enough to use the vacuum insertion approximation, by considering various ambiguities in choosing the model parameter values.) Then, the neutral Higgs contribution to the neutral D meson mass difference is given by

$$\Delta m_D(\phi) = \left| \frac{1}{m_D} \langle \bar{D}^0 | \mathcal{H}_\phi^{\Delta c=2} | D^0 \rangle \right|. \quad (21)$$

It is seen from eq.(16) that $(\Lambda_a)_{12}$ is, in particular, proportional to m_c/m_U rather than m_u/m_U , providing a significant contribution to the D^0 - $\bar{D}^0$ mixing. The Z boson contribution to the D^0 - $\bar{D}^0$ mixing is investigated in ref. [8] by considering a possible significant FCNC between the c and u quarks. In contrast to that analysis, the Z boson FCNC of u and c arises at the order of $(m_u m_c / m_U^2) \epsilon_{u-U}^2$ with the quark mass matrix (3) respecting the relation $\lambda_{u_i} \sim m_{u_i}/v$. Hence its contribution to the D^0 - $\bar{D}^0$ mixing becomes rather small in the present case.

Important contributions are also provided to the NEDM from the quark mixings with singlet quarks. Here, the u quark EDM, which is one of the main components of the NEDM, is induced by the u - U mixings in the one-loop diagrams involving the quark and neutral Higgs intermediate states. (The gauge couplings, on the other hand, do not contribute to the NEDM at the one-loop level, which is due to the same situations as in the minimal standard model [15, 10].) The total contribution of the neutral Higgs fields ϕ_a to the u quark EDM is calculated by a formula

$$d_u(\phi) = -\frac{2e}{3(4\pi)^2} \sum_a \sum_{\mathcal{U}_K=u_i, U} \left\{ \text{Im} [(\Lambda_a)_{1K} (\Lambda_a)_{K1}] \frac{m_{\mathcal{U}_K}}{m_{\phi_a}^2} I(m_{\mathcal{U}_K}^2 / m_{\phi_a}^2) \right\}, \quad (22)$$

where

$$I(X) = \frac{1}{(1-X)^2} \left[-\frac{3}{2} + \frac{1}{2}X - \frac{1}{1-X} \ln X \right], \quad (23)$$

and $m_{\mathcal{U}_K} = m_{u_i}, m_U$ represent the quark mass eigenvalues. (This form of $I(X)$ may be modified for the intermediate state of $\mathcal{U}_K = u_1$. However, the contribution with the u quark intermediate state is in any case negligible due to the very small mass m_u .) These contributions to the u quark EDM in fact arise at the first order of the u quark mass m_u ($\lambda_{u_1} \sim m_u/v$). This feature is understood by considering the limit $\lambda_{u_1} \rightarrow 0$, where the u quark EDM is vanishing due to a relevant chiral symmetry. It is also noticed from eq.(16)

that the top quark contributions with the couplings factors $(\Lambda_a)_{13}(\Lambda_a)_{31} \sim (m_u m_t / m_U^2) \epsilon_{u-U}^4$ become important together with those of the singlet U quark intermediate states.

We now make a detailed analysis of the u - U mixing effects on D^0 - $\bar{D}^0$ mixing and the u quark EDM. Numerical calculations are performed systematically in the following way: First, by taking the model parameters in certain reasonable ranges, the quark mass matrix $\mathcal{M}_U$ is diagonalized numerically to obtain the quark mixing matrices $\mathcal{V}_{U_L}$ and $\mathcal{V}_{U_R}$. Then, the quark couplings Λ_a to the neutral Higgs fields are determined with eqs. (13) and (14). Finally, $\Delta m_D(\phi)$ and $d_u(\phi)$ are calculated by using the formulae (17) – (23).

Practically, the relevant model parameters are taken as follows: The Yukawa couplings to the singlet Higgs field may be parametrized by considering eqs.(4) and (9) as

$$(f_U)_i = \kappa_i \epsilon_{u-U} e^{i\alpha_i} (|M_U|/v_S) , \quad (f'_U)_i = \kappa'_i \epsilon_{u-U} e^{i\alpha'_i} (|M_U|/v_S) , \quad (24)$$

$$\lambda_U e^{i\phi_S} + \lambda'_U e^{-i\phi_S} \equiv |\lambda_U| e^{i\beta} + |\lambda'_U| e^{i\beta'} = \sqrt{2} (M_U/v_S) . \quad (25)$$

The generation dependence of the u - U couplings is taken into account with the factors κ_i and κ'_i . The λ_U and λ'_U couplings are taken under the condition in eq. (4) so as to reproduce the given value of $|M_U|$ with varying v_S , where $|M_U|$ is approximately equal to the U quark mass m_U up to the corrections due to the u - U mixings. More specifically these parameters are taken in the ranges of

$$\epsilon_{u-U} \sim 0.1 , \quad 0 \leq \kappa_i, \kappa'_i \leq 1 , \quad 0 \leq \alpha_i, \alpha'_i < 2\pi , \quad (26)$$

$$|M_U| \simeq m_U \sim \text{several} \times 100 \text{GeV} , \quad 0 \leq \beta, \beta' < 2\pi . \quad (27)$$

Here the U quark with $m_U \sim m_t$ is favored, since significant contributions to the electroweak baryogenesis are then expected to be obtained through the CP violating t - U mixing [12]. (It is, however, necessary to take at least $m_U > m_t$, since the U quark has a dominant decay mode $U \rightarrow b + W$ similarly to the top quark.) The diagonal coupling matrix λ_u for the ordinary up-type quarks is, on the other hand, chosen suitably with the relation (10) so that the quark masses m_u , m_c and m_t are reproduced within the experimentally determined ranges [13]. (The corrections due to the u - U mixings in eq. (10) are actually at most of 10 % for $\epsilon_{u-U} \sim 0.3$.) The decay constant of the D meson is taken to be $f_D = 0.3 \text{GeV}$ [13]. The parameters concerning the Higgs fields are taken as

$$m_{\phi_0} \sim 100 \text{GeV} , \quad m_{\phi_1}, m_{\phi_2} \sim v_S \gtrsim 100 \text{GeV} , \quad (28)$$

$$\mathbf{O}_{01}, \mathbf{O}_{02} \sim v/v_S, \quad \mathbf{O}_{12} \sim 1, \quad 0 \leq \phi_S < 2\pi. \quad (29)$$

Here the mass of ϕ_0 is fixed to be a somewhat smaller value of 100 GeV or so, as suggested from the requirement that the first order electroweak phase transition be strong enough. The mixings between h_1 and s_1, s_2 are supposed to scale as v/v_S in a viewpoint of naturalness, since $\phi_0 \approx h_1$ (the standard neutral Higgs) for the extreme case of $v_S \gg v \simeq 246\text{GeV}$.

The resultant $|d_u(\phi)|$ versus $\Delta m_D(\phi)$ are shown in figs. 1 and 2 together with the experimental upper bounds on the NEDM (dashed line) and on $\Delta m_D \equiv |m_{D_1^0} - m_{D_2^0}|$ (dotted line) [13], where the relevant model parameters are taken typically as $\epsilon_{u-U} = 0.3$, $M_U = 300\text{ GeV}$ ($\simeq m_U$), $m_{\phi_0} = 100\text{ GeV}$, and $m_{\phi_1} = m_{\phi_2} = 0.5v_S$ with $v_S = 400\text{ GeV}$ (fig. 1), and 4 TeV (fig. 2). In these scatter plots, each dot corresponds to a random choice for the set of parameters such as: the complex phases $\alpha_i, \alpha'_i, \beta, \beta'$ and ϕ_S ; the generation dependent factors κ_i and $\kappa'_i = 2/3 - 1$; the coupling ratio $|\lambda'_U|/|\lambda_U| = 0.1 - 10$; and the Higgs mixing matrix $\mathbf{O}$. It is clearly found in fig. 1 that the neutral Higgs contributions are comparable to or smaller than the experimental bounds even for the case of $v_S \sim v$ with $\epsilon_{u-U} \sim 0.3$. On the other hand, if $v_S \sim 1\text{TeV}$ or larger, they become sufficiently below the experimental bounds, as seen in fig. 2. It should also be remarked that if $\kappa_{1,2}, \kappa'_{1,2} \ll 1$ while $\kappa_3, \kappa'_3 \sim 1$, namely, that only the top quark has a significant coupling to the singlet U quark, then these flavor nonconserving and CP violating effects for the light quarks become much smaller.

The modification in the CKM matrix (11) induced by the $u-U$ mixings has been estimated by the numerical calculations for the parameter choices taken in figs. 1 and 2. It is roughly given as

$$\begin{pmatrix} V_{u_L}^\dagger V_0 - V_0 \\ \epsilon_{u_L}^\dagger V_0 \end{pmatrix} \sim \begin{pmatrix} 10^{-4} & 10^{-4} & 10^{-5} \\ 10^{-4} & 10^{-4} & 10^{-4} \\ 10^{-5} & 10^{-4} & 10^{-2} \\ 10^{-3} & 10^{-2} & 10^{-1} \end{pmatrix}. \quad (30)$$

In particular, for the charged gauge interactions of the t quark and the U quark, $|V_{tb}| = 0.95 - 1$ and $|V_{Ub}| = 0 - 0.2$ are obtained. This amount of slight modification in the electroweak gauge interactions of the quarks provide contributions smaller than about 0.2 to the oblique parameters [7], which are still consistent with the experimental bounds (see for instance p. 104 in ref. [13]).

The modification in the neutral currents $\mathcal{U}^\dagger \sigma^\mu V_Z \mathcal{U}$ coupled to the Z boson has also been estimated as

$$V_Z - V_Z^{(0)} \sim \begin{pmatrix} 10^{-11} & 10^{-9} & 10^{-7} & 10^{-6} \\ 10^{-9} & 10^{-7} & 10^{-4} & 10^{-4} \\ 10^{-7} & 10^{-4} & 10^{-2} & 10^{-1} \\ 10^{-6} & 10^{-4} & 10^{-1} & 10^{-2} \end{pmatrix}, \quad (31)$$

where $V_Z^{(0)}$ represents the usual neutral currents in the absence of u - U mixings. (The neutral currents of $\mathcal{U}^c$ is not modified.) This should be compared to the quark couplings to the neutral Higgs fields, which have been estimated as

$$\Lambda_a \sim \begin{pmatrix} 10^{-5} & 10^{-4} & 10^{-2} & 10^{-1} \\ 10^{-6} & 10^{-3} & 10^{-2} & 10^{-1} \\ 10^{-6} & 10^{-4} & 10^{-1} & 10^{-1} \\ 10^{-6} & 10^{-4} & 10^{-1} & 10^{-1} \end{pmatrix}. \quad (32)$$

This indicates that the FCNC's in neutral Higgs couplings are more important than those in Z boson couplings in these sorts of models with the quark mass matrix of the form given in eq.(3) respecting the quark mass hierarchy.

As for the case of d - D mixings, the neutral Higgs contributions to the K^0 - $\bar{K}^0$ mixing and the NEDM should be investigated as well. It has actually been checked by numerical calculations that the contributions to the d quark EDM are analogous to those obtained for the u quark EDM in the presence of u - U mixings. It should also be mentioned that the CP violation parameter ϵ for the K^0 - $\bar{K}^0$ mixing, in particular, can be as large as 10^{-3} for $\epsilon_{d-D} \sim 0.1$ and $v_S \sim v$. Detailed analyses will be presented elsewhere.

Finally, some comments are presented for possible variants of the present model. (i) The complex Higgs field S may be replaced by a real field with $f'_U = \lambda'_U = 0$. Then, the coupling matrix λ_U can be made real and diagonal by a redefinition among the U_0 and U_0^c fields. Even in this case, similar contributions are obtained to Δm_D and d_u with more than one U quarks and the complex Yukawa couplings f_U . (If only one U quark is present, the complex phases in f_U are absorbed by the u_0^c fields, resulting in the vanishing of $d_u(\phi)$.) It should, however, be mentioned that with only one real singlet Higgs field the t - U mixing is ineffective for electroweak baryogenesis. This is because the complex phases in the t - U couplings to the real singlet Higgs field can be eliminated by rephasing the U_0 and U_0^c fields. (ii) It may be considered that the singlet Higgs field S is absent, regarding Δ_{u-U} and M_U as explicit mass terms in eq. (3). Even in this case, the significant FCNC's are still present in the quark

couplings to the standard neutral Higgs field $\phi_0 = h_1$. It should, however, be mentioned that the one-loop neutral Higgs contribution to d_u vanishes, just as does the one-loop Z boson contribution. This is because the standard neutral Higgs field h_1 and the Nambu-Goldstone mode h_2 couple to the quarks in the same way.

In conclusion, flavor nonconserving and CP violating effects of the quark mixings have been investigated in electroweak models incorporating singlet quarks. It is found especially that the neutral Higgs contributions to the neutral D meson mass difference and the NEDM are still consistent with the present experimental bounds even for the case where the singlet Higgs mass scale is comparable to the electroweak scale and a significant t - U mixing is present. This, in particular, implies that there is a good chance for these types of models with singlet quarks to generate a sufficient amount of baryon number asymmetry in the electroweak phase transition. A sufficiently strong electroweak phase transition can be realized with the singlet Higgs field with $v_S \sim v$, and the asymmetries of certain quantum numbers contributing to the baryon number chemical potential can be produced by the CP violating t - U mixing through the bubble wall.

We would like to thank R. N. Mohapatra and also F. del Aguila and J. A. Aguilar-Saavedra for informing us of their papers, which are relevant for the present article.

References

- [1] For a review see for instance, A. G. Cohen, D. B. Kaplan and A. E. Nelson, *Ann. Rev. Nucl. Part. Sci.* 43 (1993) 27, and references therein;
K. Funakubo, *Prog. Theor. Phys.* 46 (1996) 652, and refereces therein.
- [2] G. C. Branco and L. Lavoura, *Nucl. Phys. B* 278 (1986) 738.
- [3] F. del Aguila and J. Cortés, *Phys. Lett. B* 156 (1985) 243.
- [4] K. S. Babu and R. N. Mohapatra, *Phys. Rev. Lett.* 62 (1989) 1079.
- [5] P. Langacker and D. London, *Phys. Rev. D* 38 (1988) 886;
Y. Nir and D. Silverman, *Phys. Rev. D* 42 (1990) 1477;
L. Lavoura and J. P. Silva, *Phys. Rev. D* 47 (1993) 1117;
G. C. Branco, T. Morozumi, P. A. Parada and M. N. Rebelo, *Phys. Rev. D* 48 (1993) 1167;
V. Barger, M. S. Berger and R. J. N. Phillips, *Phys. Rev. D* 52 (1995) 1663.
- [6] L. Bento and G. C. Branco, *Phys. Lett. B* 245 (1990) 599.
- [7] L. Lavoura and J. P. Silva, *Phys. Rev. D* 47 (1993) 2046.
- [8] G. C. Branco, P.A. Parada and M.N. Rebelo, *Phys. Rev. D* 52 (1995) 4217.
- [9] W.-S. Hou and H.-c. Kao, *Phys. Lett. B* 387 (1996) 544;
G. Bhattacharyya, G. C. Branco and W.-S. Hou, *Phys. Rev. D* 54 (1996) 2114.
- [10] Y. Takeda, I. Umemura, K. Yamamoto and D. Yamazaki, *Phys. Lett. B* 386 (1996) 167.
- [11] F. del Aguila, J. A. Aguilar-Saavedra and G. C. Branco, UG-FT-69/97, hep-ph/9703410, March 1997.
- [12] J. McDonald, *Phys. Rev. D* 53 (1996) 645;
T. Uesugi, A. Sugamoto and A. Yamaguchi, *Phys. Lett. B* 392 (1997) 389.

- [13] Particle Data Group, Phys. Rev. D 54 (1996) 1.
- [14] B. McWilliams and O. Shanker, Phys. Rev. D 22 (1980) 2853.
- [15] For a review see for instance, S. M. Barr and W. J. Marciano, *CP Violation*, ed. C. Jarlskog, World Scientific, Singapore, 1989, p.455.

Figure Captions

Fig. 1 The neutral Higgs contributions of $|d_u(\phi)|$ versus $\Delta m_D(\phi)$ are plotted for the case of $v_S = 400$ GeV together with the experimental upper bounds on the NEDM (dashed line) and on Δm_D (dotted line). The relevant parameters are taken to be $\epsilon_{u-U} = 0.3$, $M_U = 300\text{GeV}$, $m_{\phi_0} = 100\text{GeV}$, and $m_{\phi_1} = m_{\phi_2} = 0.5v_S$.

Fig. 2 The neutral Higgs contributions of $|d_u(\phi)|$ versus $\Delta m_D(\phi)$ are plotted for the case of $v_S = 4$ TeV. The same values are taken for the relevant parameters as in fig. 1.




Fig. 1




Fig. 2